




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 211/28, 211/94, A01N 43/40	A1	(11) International Publication Number: WO 99/14193 (43) International Publication Date: 25 March 1999 (25.03.99)
(21) International Application Number: PCT/US98/19172 (22) International Filing Date: 14 September 1998 (14.09.98) (30) Priority Data: 60/059,150 17 September 1997 (17.09.97) US (71) Applicant: FMC CORPORATION [US/US]; 1735 Market Street, Philadelphia, PA 19103 (US). (72) Inventors: YEAGER, Walter, H.; 274 Hickory Road, Yardley, PA 19067 (US). SILVERMAN, Ian, R.; 30 East Harris Avenue, Moorestown, NJ 08057 (US). HENRIE, Robert, N., II; 785 Old York Road, East Windsor, NJ 08520 (US). CULLEN, Thomas, G.; 22 Harrison Avenue, Milltown, NJ 08850 (US). PEAKE, Clinton, J.; 29 Fran Avenue, Trenton, NJ 08628 (US). (74) Agent: CALDWELL, John, W.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: INSECTICIDAL OXIMINO AND HYDRAZONO DERIVATIVES OF N-BENZYL-4-BENZHYDRYL- AND N-BENZYL-4BENZHYDROLPIPERIDINES <div style="text-align: center;">  </div>		
(57) Abstract <p>It has now been found that certain novel oximino and hydrazone derivatives of N-benzyl-4-benzhydryl- and 4-benzhydropiperidines, and their corresponding N-oxides and agriculturally acceptable salts are useful as insecticides. These compounds are represented by formula (I) where R is hydrogen, halogen, alkyl, alkoxy, dialkylamino; R¹ is hydrogen, alkyl, haloalkyl, alkoxyalkyl, alkylcarbonyl and alkylaminocarbonyl; X is oxygen, or NR², Q is fluoro and hydroxy; Z is halogen, haloalkyle, haloalkoxy, pentahalothio, haloalkylthio, haloalkylsulfinyl, haloalkylsulfonyl, and -OCF₂O- attached to two adjacent carbon atoms of the phenyl ring; n is 0 or 1; and, when X is NR², R² is hydrogen, alkyl, alkylcarbonyl, alkoxyalkyl, or R¹ and R² taken together may be -C_mH_{2m}, or -C₂H₄OC₂H₄-, where m is 3-9. Preferred compounds include those where R is hydrogen; R¹ is alkyl; Q is hydroxy; X is oxygen; and Z is trifluoromethyl or trifluoromethoxy.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**INSECTICIDAL OXIMINO AND HYDRAZONO DERIVATIVES OF
N-BENZYL-4-BENZHYDRYL- AND
N-BENZYL-4-BENZHYDROL-PIPERIDINES**


5

Background of the Invention

This invention relates generally to insecticidal compounds and methods for their use in controlling insects. In particular, it pertains to insecticidal oximino and hydrazono derivatives of N-benzyl-4-benzhydryl- and N-benzyl-4-benzhydrolpiperidines where the oximino and hydrazono groups are substituted at the 4-position of the benzyl ring, and the corresponding N-oxides of the piperidines. This invention also pertains to the agriculturally acceptable salts of these compounds and their compositions that are useful in controlling insects in agricultural crops.

There is a continuing demand for new insecticides that are safer, more effective, and less costly. Insecticides are useful for controlling insects which may otherwise cause significant damage to crops such as wheat, corn, soybeans, potatoes, and cotton, to name a few. For crop protection, insecticides are desired that can control the insects without damaging the crops, and that have no deleterious effects to other living organisms, such as mammals.

US Patents 5,569,664 and 5,639,763 disclose a class of piperidine derivatives for use as insecticides:




in which, among others,

U is $-(CH_2)_n-$ where n is 1, 2, or 3;

Q is fluoro or hydroxy;

R is selected from a heterocycle having 5 or 6 ring atoms, optionally fused to a benzene ring, and

5



where


V, W, Y, and Z are hydrogen; and,

- 10 X is hydrogen, hydroxy, halogen, alkyl, alkoxyalkyl, alkoxy, cycloalkylalkoxy, haloalkoxy, alkenyloxy, haloalkenyloxy, alkynyloxy, alkylsilyloxy, alkylthio, haloalkylthio, cyano, cyanoalkoxy, nitro, amino, monoalkylamino, dialkylamino, alkylaminoalkoxy, alkylcarbonylamino, alkylcarbonyl, alkoxy carbonylamino, alkoxy carbonyl, alkylaminocarbonyl, aminocarbonyloxy, phenyl, phenylalkoxy, phenoxy, phenoxyalkyl, or a five-
 15 or six-membered heterocycle; each cycloalkyl moiety, heterocycle, or phenyl ring is optionally substituted with halogen, alkoxy, or haloalkoxy;
 W and X taken together may be $-OCH_2CH_2O-$, $-CH_2C(CH_3)_2O-$, $-OC(CH_3)_2O-$, $-N=C(C_2H_5)O-$, or $-CH=CHCH=C-$;
 20 R^1 and R^2 are independently selected from phenyl substituted with one or more haloalkyl or haloalkoxy.

The '664 and '763 patents do not disclose insecticidal oximino or hydrazono derivatives.

Summary of the Invention

It has now been found that certain novel oximino and hydrazono derivatives of N-benzyl-4-benzhydryl- and 4-benzhydropiperidines, and their corresponding N-oxides and agriculturally acceptable salts are useful as insecticides. These compounds are represented by Formula I:




I

where R, R¹, X, and Z are substituents described below. Preferred compounds include those where R is hydrogen; R¹ is alkyl; Q is hydroxy; X is oxy; and Z is trifluoromethyl or trifluoromethoxy.

Detailed Description of the Invention

Certain novel oximino and hydrazono derivatives of N-benzyl-4-benzhydryl- and 4-benzhydropiperidines, their corresponding N-oxides and their agriculturally acceptable salts are useful in controlling insects. These compounds are represented by formula I:



I

where

R is hydrogen, halogen, alkyl, alkoxy, dialkylamino;

R¹ is hydrogen, alkyl, haloalkyl, alkoxyalkyl, alkylcarbonyl and
alkylaminocarbonyl;

5 Q is fluoro or hydroxy;

X is oxygen or NR²;

Z is halogen, haloalkyl, haloalkoxy, pentahalothio, haloalkylthio,
haloalkylsulfinyl, haloalkylsulfonyl, and -OCF₂O-attached to two adjacent
carbon atoms of the phenyl ring;

10 n is 0 or 1; and,

when X is NR², R² is hydrogen, alkyl, alkylcarbonyl, alkoxy carbonyl, or R¹ and
R² taken together is -C_mH_{2m}-, - or -C₂H₄OC₂H₄-, where m is 3-9.

Preferred compounds are those where R is hydrogen; R¹ is alkyl; Q is
15 hydroxy; X is oxygen; Z is trifluoromethyl or trifluoromethoxy; and n is 0 or 1.


A particularly preferred compound is
N-[4-(ethoxyiminomethyl)phenylmethyl]-4-[bis(4-trifluoromethoxyphenyl)hydro
xymethyl]piperidine N-oxide.

20

As used in this specification and, unless otherwise indicated, the term
"alkyl" and "alkoxy" used alone or as part of a larger moiety includes 1 to 6
carbon atoms either straight or branched chain. "Halogen" refers to chlorine,
bromine, and fluorine. The term "TEA" refers to triethylamine, "DMSO" refers
25 to dimethyl sulfoxide, "DMF" refers to N,N-dimethylformamide, "DAST" refers
to diethylaminosulfur trifluoride, and "NCS" refers to N-chlorosuccinimide.


The oximino and hydrazono derivatives of Formula I may be prepared
by methods described below or by methods known in the art for similar
30 compounds.

Scheme 1



Scheme 1 shows one approach to preparing compounds of the
 present invention where R is hydrogen or alkyl. In general,
 N-benzyl-4-benzhydryl- or N-benzyl-4-benzhydropiperidines may be prepared
 as described in US Patent 5,569,664 by reacting a 4-benzhydryl- or a
 4-benzhydropiperidine with a benzyl halide. In an analogous manner, a
 benzyl halide having in the para position a carboxaldehyde or keto group may
 be treated with a 4-benzhydryl- or a 4-benzhydropiperidine to provide
 intermediate 2. For example, 4-bromomethylbenzaldehyde or
 4'-(bromomethyl)acetophenone may be treated with
 4-[bis(4-trifluoromethoxyphenyl)hydroxymethyl]piperidine under basic
 conditions in a suitable solvent such as DMSO, affording 2 where Z is
 trifluoromethoxy and R is formyl or acetyl. Intermediate 2 may then be
 condensed with a hydrazine or alkoxyamine, such as methoxylamine
 hydrochloride or 1,1-dimethylhydrazine, to provide I.

6

Scheme 2

(a) NCS, HCl (gas), DMF (b) $\text{HN(CH}_3)_2$, THF (c) $n\text{-BuLi}$, DMF, THF
 (d) NaBH_4 , $\text{C}_2\text{H}_5\text{OH}$ (e) SOCl_2 , Pyridine

5 Scheme 2 shows another approach to compounds of the present invention where the oximino or hydrazono groups are introduced earlier in the synthesis. As an example, an oxime ether of 4-bromobenzaldehyde 3 may be chlorinated as shown in step (a) and then treated with a dialkylamine according to step (b) to provide an amidine ether intermediate 4. The bromine of 4 may be replaced by a chloromethyl group in three steps: (c) adding DMF to a lithiated species to provide the corresponding benzaldehyde, (d) reduction to the benzyl alcohol, and (e) chlorination to the benzyl chloride 5. Benzylation of a 4-benzhydropiperidine with benzyl chlorides 5 as described above provide further compounds of the present invention.

20 The N-benzyl-4-benzhydropiperidines I or 2 (Q is hydroxy) may be further transformed into the benzhydryl fluorides I (Q is fluoro) using either DAST or HF-pyridine.

The n-oxides I ($n=1$) are prepared by the treatment of the parent

piperidine compound I (n=0) with 30% hydrogen peroxide in a suitable solvent. In addition the agriculturally acceptable salts are also prepared by methods similar to those known in the art for similar compounds.

- 5 A procedure for a method useful to prepare compounds of this invention is given in the example below.

EXAMPLE 1

Synthesis of
N-[4-(ethoxyiminomethyl)phenylmethyl]-4-[bis(4-trifluoromethoxyphenyl)hydroxymethyl]piperidine
(Compound 15)

15 A solution of 1.0 gram (0.0018 mole) of
N-[4-formylphenylmethyl]-4-[bis(4-trifluoromethoxyphenyl)hydroxymethyl]piper
idine, 1.8 grams (0.018 mole) of O-ethylhydroxylamine hydrochloride, and 2.5
mL (0.018 mole) of TEA in 50 mL of ethanol was stirred at ambient
temperature for about four hours. The reaction mixture was then
20 concentrated under reduced pressure to near-dryness. The residue was
taken up in 100 mL of water and extracted with three 75 mL portions of
methylene chloride. The combined extracts were washed with 75 mL of
aqueous 10% lithium chloride solution and dried with sodium sulfate. The
mixture was filtered and concentrated under reduced pressure to a residue.
25 The residue was purified using column chromatography on silica gel, eluting
with ethyl acetate-chloroform mixtures. The product-containing fractions were
combined and concentrated under reduced pressure, yielding 0.6 gram of
Compound 15, mp 46-49° C. The NMR spectrum was consistent with the
proposed structure. The reaction was repeated several times on a larger
30 scale.


EXAMPLE 2

Synthesis of
N-[4-(ethoxyiminomethyl)phenylmethyl]-4-[bis(4-trifluoromethoxyphenyl)hydro-
xymethyl]piperidine N-oxide
(Compound 16)

A solution of 22.7 grams (0.030 mole) of
N-[4-(ethoxyiminomethyl)phenylmethyl]-4-[bis(4-trifluoromethoxyphenyl)hydro-
xymethyl]piperidine and 50 mL of 30% hydrogen peroxide in 300 mL of
methanol was stirred at ambient temperature for about 16 hours. After this
time the reaction mixture was diluted with 500 mL of water and extracted with
three 200 mL portions of ethyl acetate. The combined extracts were washed
with 200 mL of aqueous 10% lithium chloride. The organic layer was dried
with sodium sulfate and filtered. The filtrate was concentrated under reduced
pressure to a residue. The residue was purified using column
chromatography on silica gel, eluting with methanol-methylene chloride
mixtures. The product-containing fractions were combined and concentrated
under reduced pressure, yielding 22.1 grams of Compound 16, mp 188-191°
C. The NMR spectrum was consistent with the proposed structure.

Table 1 below shows representative compounds of the present
invention.

Table 1



where Q is hydroxy

	Cmpd No	R	R ¹	n	X	Z
5						
10	1	H	H	0	O	4-CF ₃
	2	H	H	1	O	4-CF ₃
15	3	H	CH ₃	0	O	4-CF ₃
	4	H	CH ₃	1 monohydrate	O	4-CF ₃
	5	H	C ₂ H ₅	0	O	4-CF ₃
20	6	H	C ₂ H ₅	1	O	4-CF ₃
	7	H	n-C ₃ H ₇	0	O	4-CF ₃
25	8	H	i-C ₃ H ₇	1	O	4-CF ₃
	9	H	C ₂ H ₄ OCH ₃	0	O	4-CF ₃
	10	OCH ₃	C ₂ H ₅	0	O	4-CF ₃
30	11	N(CH ₃) ₂	CH ₃	0	O	4-CF ₃

10

	12	$N(CH_3)_2$	CH_3	1	O	4- CF_3
	13	H	H	0	O	4- OCF_3
5	14	H	CH_3	0	O	4- OCF_3
	15	H	C_2H_5	0	O	4- OCF_3
10	16	H	C_2H_5	1	O	4- OCF_3
	17	H	i- C_3H_7	0	O	4- OCF_3
	18	H	n- C_3H_7	1	O	4- OCF_3
15	19	H	$C_3H_6OCH_3$	1	O	4- OCF_3
	20	CH_3	H	0	O	4- OCF_3
20	21	CH_3	CH_3	0	O	4- OCF_3
	22	CH_3	C_2H_5	0	O	4- OCF_3
	23	OC_2H_5	C_2H_5	0	O	4- OCF_3
25	24	H	C_2H_5	0	O	4- SF_5
	25	H	C_2H_5	1	O	4- SF_5

30 where Q is hydroxy, X is NR^2

	Cmpd No	R	R^1	R^2	n	Z
35	26	CH_3	$-C_2H_4OC_2H_4-$		0	4- CF_3
	27	H	CH_2CF_3	H	0	4- OCF_3

11

5	28	H	CH ₃	CH ₃	0	4-OCF ₃
	29	CH ₃	CH ₃	CH ₃	0	4-OCF ₃
	30	H		-C ₄ H ₈ -	0	4-OCF ₃
	31	CH ₃		-C ₄ H ₈ -	0	4-OCF ₃
	32	H		-C ₂ H ₄ OC ₂ H ₄ -	0	4-OCF ₃
10	33	H	CH ₃	CH ₃	0	4-SF ₅
	34	CH ₃		-C ₂ H ₄ OC ₂ H ₄ -	0	4-OCF ₃
15	35	H	C ₄ H ₉	CO ₂ CH ₃	0	4-CF ₃
	36	H	C ₄ H ₉	COCH ₃	0	4-CF ₃
20	37	H	C ₄ H ₉	CO ₂ CH ₃	0	4-OCF ₃
	38	H	C ₄ H ₉	COCH ₃	0	4-OCF ₃

where Q is Hydroxy

25

	Cmpd No	R	R ¹	n	X	Z
30	39	H	CH ₃	1 ethanesulfon ic acid salt	O	4-CF ₃
	40	H	C ₂ H ₅	0 HCl salt	O	4-CF ₃
	41	H	i-C ₃ H ₇	0	O	4-CF ₃

	42	H	n-C ₃ H ₇	1	O	4-CF ₃
	43	H	CH ₂ CH ₂ F	0	O	4-CF ₃
5	44	H	CONHC ₂ H ₅	0	O	4-CF ₃
	45	H	COCH(CH ₃) ₂	0	O	4-CF ₃
10	46	H	CH ₂ CH ₂ F	1	O	4-CF ₃
	47	H	COCH(CH ₃) ₂	1	O	4-CF ₃
	48	H	COCH ₃	0	O	4-CF ₃
15	49	H	CONHCH ₃	0	O	4-CF ₃
	50	Cl	C ₂ H ₅	0	O	4-CF ₃
20	51	H	C ₂ H ₅	1 ethanesulfonic acid salt	O	4-CF ₃
	52	H	CONHCH ₃	1	O	4-CF ₃
	53	H	COCH(CH ₃) ₂	0	O	4-OCF ₃
25	54	H	COCH ₃	0	O	4-OCF ₃
	55	H	CH ₂ CH ₂ F	0	O	4-OCF ₃
	56	H	CONHC ₂ H ₅	0	O	4-OCF ₃
30	57	H	CH ₃	1	O	4-OCF ₃
	58	H	CH ₃	0	O	4-SF ₅
35	59	H	CH ₃	1	O	4-SF ₅

5	60	H	CH ₃	0	O	3-OCF ₂ O-4
	61	H	CH ₃	1	O	3-OCF ₂ O-4
	62	H	CH ₃	0	O	4-SCF ₃
	63	H	CH ₃	1	O	4-SCF ₃
10	64	H	C ₂ H ₅	0	O	3-OCF ₂ O-4
	65	H	C ₂ H ₅	1	O	3-OCF ₂ O-4
	66	H	C ₂ H ₅	0	O	4-SCF ₃
15	67	H	C ₂ H ₅	1	O	4-SCF ₃
	68	H	CH ₃	0	O	4-SOCF ₃
20	69	H	CH ₃	1	O	4-SOCF ₃
	70	H	CH ₃	0	O	4-SO ₂ CF ₃
	71	H	CH ₃	1	O	4-SO ₂ CF ₃
25	72	H	C ₂ H ₅	0	O	4-SOCF ₃
	73	H	C ₂ H ₅	1	O	4-SOCF ₃
30	74	H	C ₂ H ₅	0	O	4-SO ₂ CF ₃
	75	H	C ₂ H ₅	1	O	4-SO ₂ CF ₃

where Q is fluoro

35

Cmpd No	R	R ¹	n	X	Z
---------	---	----------------	---	---	---

14

5	76	H	CH ₃	0	O	4-CF ₃
	77	H	CH ₃	1	O	4-CF ₃
	78	H	CH ₃	0	O	4-OCF ₃
	79	H	CH ₃	1	O	4-OCF ₃
10	80	H	C ₂ H ₅	0	O	4-CF ₃
	81	H	C ₂ H ₅	1	O	4-CF ₃
	82	H	C ₂ H ₅	0	O	4-OCF ₃
15	83	H	C ₂ H ₅	1	O	4-OCF ₃

Table 2
Characterizing Data

20	Cmpd. No.	MP (C°)/ Physical State	Cmpd. No.	MP (C°)/ Physical State	Cmpd. No.	MP (C°)/ Physical State
25	1	108-117	2	217-221 dec	3	65-71
	4	195-203	5	60-65	6	207-210 dec
	7	50-59	8	201-205	11	66-68
30	12	165-168	13	91-97	14	47-52
	15	46-49	16	188-191	20	87-94
	21	56-62	22	56-62	27	56-62
35						

	28	58-65	29	51-55	30	69-72
	31	49-58	32	75-81	34	63-67
5	41	85-88	42	200-205	43	63-68
	44	178-179	45	78-87	46	203-205
10	47	130-140	48	78-87	49	98-105
	50	Oil	51	202-209	52	184-194
	53	65-70	54	73-82	55	Oil
15	56	138-145				

Biological Testing

20 Candidate insecticides were evaluated for activity against tobacco budworm (Heliothis virescens [Fabricius]) by applications to the surface of a wheat germ-based artificial insect diet. Solutions of the candidate insecticides were prepared for testing by diluting a standard 50 millimolar DMSO solution of each candidate insecticide with DMSO, then further

25 diluting with a 1:1 water/acetone solution (V/V). Forty microliters of this solution of calculated concentration was then pipetted onto the surface of the diet in each of six containers, to provide six replicates for each rate of application. Once treated, the contents of the containers were allowed to dry, leaving the calculated concentration (in millimoles) of candidate

30 insecticide on the surface of the diet. In each container was placed one second instar tobacco budworm larvae. The container was sealed with a transparent film, and then held in a growth chamber for five days. After the five-day exposure period the insecticidal activity of the candidate insecticide was recorded as percent mortality when compared to the total number of

insects infested.

The compounds of the present invention were tested in the laboratory as solutions in DMSO. It is expected that all formulations normally employed in applications of insecticides would be usable with the compounds of the present invention. These include wettable powders, emulsifiable concentrates, water suspensions, flowable concentrates, and the like.

Insecticidal activity at a selected molar concentration of test chemical are given for various compounds of this invention in Table 3. The test compounds are identified by numbers which correspond to those in Table 1.

Table 3
Insecticidal Activity of Test Compounds Applied to the Surface of the Diet of Tobacco Budworm

	Compound No.	1	2	3	4	5	6	7	8	11
20	Percent Mortality	100	67	100	100	100	100	100	100	67
25	Compound No.	12	13	14	15	16	20	21	22	27
	Percent Mortality	100	100	100	100	100	100	100	100	100
30	Compound No.	28	29	30	31	32	34	41	42	43
35	Percent Mortality	100	17	100	0	50	67	100	50*	100

17

5	Compound No.	44	45	46	47	48	49	50	51	52
	Percent Mortality	83	100	100	100	67	50	50	100	83

10	Compound No.	53	54	55	56
	Percent Mortality	100	100	100	83

15 Unless otherwise noted the concentration of candidate insecticide on the surface of the diet is 0.025 millimoles

*Concentration of candidate insecticide on the surface of the diet is 0.0025 millimoles

20 For insecticidal application, the active compounds are formulated into insecticidal compositions by admixture in insecticidally effective amount with adjuvants and carriers normally employed in the art for facilitating the dispersion of active ingredients for the particular utility desired, recognizing the fact that the formulation and mode of application of a toxicant may affect the activity of the material in a given application. Thus, for agricultural use the present insecticidal compounds may be formulated as granules of relatively large particle size, as water-soluble or water-dispersible granules, as powdery dusts, as wettable powders, as emulsifiable concentrates, as solutions, or as any of several other known types of formulations, depending on the desired mode of application.

35 These insecticidal compositions may be applied either as water-diluted sprays, or dusts, or granules to the areas in which insect control is desired. These formulations may contain as little as 0.1%, 0.2% or 0.5% to as much as 95% or more by weight of active ingredient.

Dusts are free flowing admixtures of the active ingredients with finely divided solids such as talc, natural clays, kieselguhr, flours such as walnut shell and cottonseed flours, and other organic and inorganic solids which act as dispersants and carriers for the toxicant; these finely divided solids have an average particle size of less than about 50 microns. A typical dust formulation useful herein is one containing 1.0 part or less of the insecticidal compound and 99.0 parts of talc.

Wettable powders are in the form of finely divided particles which disperse readily in water or other dispersant. The wettable powder is ultimately applied to the locus where insect control is desired either as a dry dust or as an emulsion in water or other liquid. Typical carriers for wettable powders include Fuller's earth, kaolin clays, silicas, and other highly absorbent, readily wet, inorganic diluents. Wettable powders normally are prepared to contain about 5-80% of active ingredient, depending on the absorbency of the carrier, and usually also contain a small amount of a wetting, dispersing, or emulsifying agent to facilitate dispersion. For example, a useful wettable powder formulation contains 80.8 parts of the insecticidal compound, 17.9 parts of Palmetto clay, and 1.0 part of sodium lignosulfonate and 0.3 part of sulfonated aliphatic polyester as wetting agents.

Other useful formulations for insecticidal applications are emulsifiable concentrates (ECs) which are homogeneous liquid compositions dispersible in water or other dispersant, and may consist entirely of the insecticidal compound and a liquid or solid emulsifying agent, or may also contain a liquid carrier, such as xylene, heavy aromatic naphthas, isophchrone, or other non-volatile organic solvent. For insecticidal application these concentrates are dispersed in water or other liquid carrier, and normally applied as a spray to the area to be treated. The percentage by weight of the essential active ingredient may vary according to the manner in which the composition is to be applied, but in general comprises 0.5 to 95% of active ingredient by weight of

the insecticidal composition.

Flowable formulations are similar to ECs except that the active ingredient is suspended in a liquid carrier, generally water. Flowables, like
5 ECs, may include a small amount of a surfactant, and contain active ingredient in the range of 0.5 to 95%, frequently from 10 to 50%, by weight of the composition. For application, flowables may be diluted in water or other liquid vehicle, and are normally applied as a spray to the area to be treated.

10 Typical wetting, dispersing, or emulsifying agents used in agricultural formulations include, but are not limited to, the alkyl and alkylaryl sulfonates and sulfates and their sodium salts; alkylaryl polyether alcohols; sulfated higher alcohols; polyethylene oxides; sulfonated animal and vegetable oils; sulfonated petroleum oils; fatty acid esters of polyhydric alcohols and the
15 ethylene oxide addition products of such esters; and the addition product of long-chain mercaptans and ethylene oxide. Many other types of useful surface-active agents are available in commerce. The surface-active agents, when used, normally comprise from 1 to 15% by weight of the composition.

20 Other useful formulations include suspensions of the active ingredient in a relatively non-volatile solvent such as water, corn oil, kerosene, propylene glycol, or other suitable solvents.

Still other useful formulations for insecticidal applications include
25 simple solutions of the active ingredient in a solvent in which it is completely soluble at the desired concentration, such as acetone, alkylated naphthalenes, xylene, or other organic solvents. Granular formulations, wherein the toxicant is carried on relatively coarse particles, are of particular utility for aerial distribution or for penetration of cover crop canopy.
30 Pressurized sprays, typically aerosols wherein the active ingredient is dispersed in finely divided form as a result of vaporization of a low boiling

dispersant solvent carrier, such as carbon dioxide, propane, or butane, may also be used. Water-soluble or water-dispersible granules are also useful formulations for insecticidal application of the present compounds. Such granular formulations are free-flowing, non-dusty, and readily water-soluble or water-miscible. The soluble or dispersible granular formulations described in U.S. Patent 3,920,442 are useful herein with the present insecticidal compounds. In use by the farmer on the field, the granular formulations, emulsifiable concentrates, flowable concentrates, solutions, etc., may be diluted with water to give a concentration of active ingredient in the range of say 0.1% or 0.2% to 1.5% or 2%.


The active insecticidal compounds of this invention may be formulated and/or applied with other insecticides, fungicides, nematicides, plant growth regulators, fertilizers, or other agricultural chemicals. In using an active compound of this invention, whether formulated alone or with other agricultural chemicals, to control insects, an effective amount and concentration of the active compound is applied to the locus where control is desired. The locus may be, e.g., the insects themselves, plants upon which the insects feed, or the insect habitat. When the locus is the soil, e.g., soil in which agricultural crops have been or will be planted, the composition of the active compound may be applied to and optionally incorporated into the soil. For most applications the effective amount may be as low as, e.g. about 10 to 500 g/ha, preferably about 100 to 250 g/ha.

It is apparent that various modifications may be made in the formulation and application of the compounds of this invention without departing from the inventive concepts herein as defined in the claims.

We claim:

1. A compound having the formula

5



I

where

- 10 R is hydrogen, halogen, alkyl, alkoxy, dialkylamino;
 R¹ is hydrogen, alkyl, haloalkyl, alkoxyalkyl, alkylcarbonyl and
 alkylaminocarbonyl;
 Q is fluoro or hydroxy;
 X is oxygen or NR²
 15 Z is halogen, haloalkyl, haloalkoxy, pentahalothio, haloalkylthio,
 haloalkylsulfinyl, haloalkylsulfonyl, and -OCF₂O-attached to two
 adjacent carbon atoms of the phenyl ring;
 n is 0 or 1;
 when X is NR², R² is hydrogen, alkyl, alkylcarbonyl, alkoxy carbonyl, or
 20 R¹ and R² taken together may be -C_mH_{2m}-, - or -C₂H₄OC₂H₄-, where m
 is 3-9; and their agriculturally acceptable salts.

25

2. A compound of claim 1 where R is hydrogen; R¹ is alkyl; Q is
 hydroxy; X is oxygen; Z is trifluoromethyl or trifluoromethoxy; n is 0 or
 1; and their agriculturally acceptable salts.
3. A compound of claim 2 which is
 N-[4-(ethoxyiminomethyl)phenylmethyl]-4-[bis(4-trifluoromethoxyphenyl)]

hydroxymethyl]piperidine N-oxide; and its agriculturally acceptable salts.

4. An insecticidal composition comprising an insecticidally effective amount of a compound of claim 1, and an insecticidally compatible carrier.
5. An insecticidal composition comprising an insecticidally effective amount of a compound of claim 2, and an insecticidally compatible carrier.
6. An insecticidal composition comprising an insecticidally effective amount of the compound of claim 3, and an insecticidally compatible carrier.
7. A method of controlling insects, comprising application to the locus where insect control is desired, an insecticidally effective amount of a composition of claim 4.
8. A method of controlling insects, comprising application to the locus where insect control is desired, an insecticidally effective amount of a composition of claim 5.
9. A method of controlling insects, comprising application to the locus where insect control is desired, an insecticidally effective amount of a composition of claim 6.

INTERNATIONAL SEARCH REPORT

In .tional Application No

PCT/US 98/19172

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D211/28 C07D211/94 A01N43/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 639 763 A (I. R. SILVERMAN ET AL.) 17 June 1997 cited in the application see claims 1-9; tables 3,4 ---	1-9
Y	US 5 569 664 A (I. R. SILVERMAN ET AL.) 29 October 1996 cited in the application see claims 1-31; tables 3,4 ---	1-9
Y	WO 96 36228 A (I. R. SILVERMAN ET AL.) 21 November 1996 see claims 1-19; tables 3,4 ---	1-9
Y	WO 95 23507 A (I. R. SILVERMAN ET AL.) 8 September 1995 see claims 1-15; tables 3,4 -----	1-9



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 December 1998

Date of mailing of the international search report

18/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herz, C

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. l. Application No

PCT/US 98/19172

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5639763 A	17-06-1997	US 5569664 A	29-10-1996
		AU 694192 B	16-07-1998
		AU 1935195 A	18-09-1995
		BG 100815 A	31-10-1997
		BR 9506972 A	23-09-1997
		CA 2184576 A	08-09-1995
		CN 1147193 A	09-04-1997
		CZ 9602564 A	12-02-1997
		EP 0748159 A	18-12-1996
		HU 75053 A	28-03-1997
		JP 9505080 T	20-05-1997
		LV 11809 A	20-08-1997
		LV 11809 B	20-12-1997
		NO 963641 A	30-10-1996
		PL 316082 A	23-12-1996
		SK 112596 A	05-03-1997
		WO 9523507 A	08-09-1995
US 5569664 A	29-10-1996	US 5639763 A	17-06-1997
WO 9636228 A	21-11-1996	AU 5753696 A	29-11-1996
		CN 1184405 A	10-06-1998
		EP 0844825 A	03-06-1998
WO 9523507 A	08-09-1995	AU 694192 B	16-07-1998
		AU 1935195 A	18-09-1995
		BG 100815 A	31-10-1997
		BR 9506972 A	23-09-1997
		CA 2184576 A	08-09-1995
		CN 1147193 A	09-04-1997
		CZ 9602564 A	12-02-1997
		EP 0748159 A	18-12-1996
		HU 75053 A	28-03-1997
		JP 9505080 T	20-05-1997
		LV 11809 A	20-08-1997
		LV 11809 B	20-12-1997
		NO 963641 A	30-10-1996
		PL 316082 A	23-12-1996
		SK 112596 A	05-03-1997
		US 5639763 A	17-06-1997